

CHARACTERISTICS OF ROOTSTOCKS

ROOTSTOCK		VIGOR	AFFINITY WITH VITIS VINIFERA	RESISTANCE TO ACTIVE LIMESTONE %	ADAPTABILITY TO THE SOIL									RESISTANCE TO PHYLLOXERA	RESISTANCE TO NEMATODES
					ACID	CLAY	DRY	FERTILI	FRESH	SANDY	SALINE	DROUGHT	HUMID		
BERLANDIERI x RIPARIA	KOBER 5 BB	****	*****	18	***	***	*	***	*****	**	*	*	***	****	****
	SO4	****	***	18	***	***	***	***	***	***	**	**	**	****	****
	BINOVA	****	***	18	***	***	*	***	*****	***	*	*	**	****	****
	420 A	***	**	15	**	***	*****	**	**	*	*	***	*	***	*
	161.49	*	**	25	**	**	*	**	**	*	*	**	*	****	*
	TELEKI 5C	***	***	12	**	*	*	***	*****	*	*	*	***	****	****
BERLANDIERI x RUPESTRIS	1103 PAULSEN	*****	*****	22	***	***	*****	*	*	****	***	****	**	*****	***
	110 RICHTER	***	*	17	**	***	*****	*	*	***	*	****	*	*****	***
	140 RUGGERI	****	*	35	*	***	*****	*	*	***	*	*****	*	*****	***
RIPARIA x RUPESTRIS	101.14	*	**	9	*	*	*	****	***	*	*	*	***	****	***
	3309 C	**	**	10	*	**	*	****	*****	*	*	*	***	****	*
	SCHWARZMANN	**	**	10	*	**	*	****	***	*	*	*	***	****	**
RIPARIA x	BÖRNER	***	***	7	*	**	****	*	*	***	*	****	*	****	****
VINIFERA x BERLANDIERI	196.17	****	**	6	***	***	**	***	****	**	***	***	*	**	*
	GRAVESAC	***	**	6	****	*	***	***	****	***	*	*	***	****	*
	41 B	***	*	35	***	*	***	***	*****	*	*	*	*	***	*
	FERCAL	***	*	35	****	**	***	***	****	*	**	***	***	****	***

***** EXCELLENT

**** GOOD

*** DISCREET

** SUFFICIENT

* INSUFFICIENT